

Wellness Center – Health Services 220 Pawtucket Street, Suite 300 Lowell, Massachusetts 01854-5144 Tel: 978-934-6800

TUBERCULOSIS RISK ASSESSMENT

(Part 1 of 2)

(PART 1 MUST BE COMPLETED ON YOUR PORTAL)

Name:	Date:				
Name:	Date: Middle Initial Phone Number:				
Home Address:					
Local Address:					
Country of Birth:					
Please answer the following questions:					
Have you ever been in close contact with anyone sick with tuberculosis?			No		
Were you born in one of the countries listed below? If yes, what country:	Date of entry to United States:	Yes	No		
Have you lived or traveled for more than one month in on If yes, what country/ies:		Yes	No		
If the answer is NO to all of the above questions, no furt	her testing is required. Submit form to Health	Services.	ı		
If the answer is YES to any of the above questions, UMASS Lowell Health Services requires that you have a tuberculosis skin test (TST) or an IGRA. The TST or IGRA must be completed no more than 6 months before the start of classes. TST preferred to be completed in USA. Continue to Part 2 of this form.					
If you have had a positive TST or IGRA in the past, you	will not need another TST. Continue to Part	2 of this for	m.		
*COUNTRIES WITH HIGH PATES OF TUREPOUT OSIS					

*COUNTRIES WITH HIGH RATES OF TUBERCULOSIS

Afghanistan	Central African Republic	Greenland	Madagascar	Pakistan	Suriname
Algeria	Chad	Guam	Malawi	Palau	Tajikistan
Angola	China	Guatemala	Malaysia	Panama	Tanzania-UR
Anguilla	China (Hong Kong SAR)	Guinea	Maldives	Papua New Guinea	Thailand
Argentina	China (Macao SAR)	Guinea-Bissau	Mali	Paraguay	Timor-Leste
Armenia	Colombia	Guyana	Marshall Islands	Peru	Togo
Azerbaijan	Congo	Haiti	Mauritania	Philippines	Tunisia
Bangladesh	Congo DR	Honduras	Mexico	Portugal	Turkmenistan
Belarus	Cote d'Ivoire	India	Micronesia	Qatar	Tuvalu
Belize	Djibouti	Indonesia	Moldova-Rep	Romania	Uganda
Benin	Dominican Republic	Iraq	Mongolia	Russian Federation	Ukraine
Bhutan	Ecuador	Kazakhstan	Morocco	Rwanda	Uruguay
Bolivia	El Salvador	Kenya	Mozambique	Sao Tome & Principe	Uzbekistan
Bosnia & Herzegovina	Equatorial Guinea	Kiribati	Myanmar	Senegal	Vanuatu
Botswana	Eritrea	Korea-DPR	Namibia	Sierra Leone	Venezuela
Brazil	Eswatini	Korea-Rep	Nauru	Singapore	Viet Nam
Brunei Darussalam	Ethiopia	Kyrgyzstan	Nepal	Solomon Islands	Yemen
Burkina Faso	Fiji	Lao PDR	Nicaragua	Somalia	Zambia
Burundi	Gabon	Lesotho	Niger	South Africa	Zimbabwe
Cabo Verde	Gambia	Liberia	Nigeria	South Sudan	
Cambodia	Georgia	Libya	Niue	Sri Lanka	
Cameroon	Ghana	Lithuania	Northern Mariana Islands	Sudan	

Rev. 06/24

^{*} Source: World Health Organization Global Health Observatory, Tuberculosis Incidence. Countries with average incidence rates of ≥ 20 cases per 100,000 population.



Wellness Center – Health Services 220 Pawtucket Street, Suite 300 Lowell, Massachusetts 01854-5144 Tel: 978-934-6800

TUBERCULOSIS RISK ASSESSMENT

(Part 2 of 2)

Name:	Date:/_ /					
Name: Last First	Date:// Middle Initial					
DOB:/SID#:	Phone Number:					
Country of Birth:						
Tuberculin Skin Test (TST) or IGRA						
Plant date: / / Read date*: /	_/ Result**:mm of induration ^Positive Negative					
	th lab report. Rapid Serology TB test not accepted.): ^Positive or Negative					
	es below. ^ If positive, continue below.					
II POSITIVE Tubercullii Skill Test of IGRA (I	now or by history) the following are required:					
Date of positive TST or IGRA:// Result: mm of induration						
Chest X-ray: Date:// Result: Normal Abnormal (must attach chest x-ray report, NOT the X-ray)						
Clinical Evaluation:NormalAbnormal						
Describe:						
Treatment:NoYes						
Drug/s, dose, frequency, and dates:						
Health Care Provider						
Name:	Address:					
Signature:	Phone:					
** Interpretation Guidelines						

* Interpretation Guidelines

5 mm or greater is positive:

- Recent close contacts of an individual with infectious TB
- Persons with fibrotic changes on a prior chest x-ray consistent with past TB disease
- Organ transplant recipients
- Immunosuppressed persons: taking >15 mg/d of prednisone for > 1 month; taking a TNF-α antagonist
- Persons with HIV/AIDS

10 mm or greater is positive:

- Persons born in a high prevalence country or who resided in one for a significant amount of time
- · History of illicit drug use
- Mycobacteriology laboratory personnel
- History of resident, worker or volunteer in high-risk congregate settings
- Persons with the following clinical conditions: silicosis, diabetes mellitus, chronic renal failure, leukemias and lymphomas, head, neck or lung cancer, low body weight (>10% below ideal), gastrectomy or intestinal bypass, chronic malabsorption syndromes

15 mm or greater is positive:

 Persons with no known risk factors for TB disease

Rev. 06/24